

Five Year Integrated M.Sc. Examination, 2019

Semester - IV

Course: PH-2-4-2 (Old) (Statistical Mechanics)

Time: Three Hours

Full Marks: 40

Questions are of value as indicated in the margin.
Answer **Question No.1** and **any three** from the rest.

1. Answer **any five** from the following: 5 x 2 = 10
 - a) Define phase space? How do you represent a microstate in phase space?
 - b) Draw the phase space orbit of a particle with energy E , confined within a one dimensional wire extended from $-L$ to L .
 - c) State equipartition theorem.
 - d) Using the definition of microcanonical partition function show that the entropy is an additive quantity for weakly interacting systems.
 - e) Write the two particle wave-function for Fermions and Bosons.
 - f) Briefly explain Gibbs' paradox with an example.
 - g) Explain Pauli's exclusion principle.
2.
 - a) Obtain the canonical probability distribution function for a system which is in thermal equilibrium with a reservoir at a temperature T K.
 - b) Obtain the expression of average energy (E), pressure (p) and entropy (S) for the canonical ensemble. 4+(2+2+2)=10
3.
 - a) State the postulate of equal a priori probabilities. Is it valid for all types of ensembles?
 - b) How does the total number of microstates for a isolated system behave at equilibrium? Define reversible and irreversible processes with examples.
 - c) Can you draw the phase space trajectory of a quantum system? Give justification in support of your answer.
 - d) Define grand canonical ensemble. Write down the expression for corresponding partition function. (2+1)+(1+2)+2+(1+1)=10
4.
 - a) Calculate the canonical partition function of an classical ideal gas consists of N number of gas particles confined within a volume V . Hence calculate the entropy and average energy of the gas.
 - b) Derive the expression for thermal wave length ($\bar{\lambda}$). How is it related with inter particle distance (R) for a quantum system. (3+2+1)+(3+1)=10
5.
 - a) Consider an isolated system in equilibrium. Establish the relation between the temperatures of two different parts of this system.
 - b) Consider an isolated system of N non-interacting particles, each fixed in position. Each particle may exist in one of the two states $E=0$ or $E= \epsilon$. Treat the particles as distinguishable.
 - i) Calculate the number of accessible states, $\Omega(E, N)$ for this system.
 - ii) Express entropy S , in terms of $\Omega(E, N)$. Apply Starling approximation to simplify this result. ($\ln N! = N \ln N - N$, for $N \rightarrow \infty$)
 - iii) Calculate the maximum value of the entropy. 4+(2+2+2)=10

P.T.O.

6. a) Consider a microcanonical ensemble of a system of N non-interacting **Fermions** confined to a volume V and sharing a total energy E . Let us assume that the energy spectrum is discrete in which the i -th level corresponds to the energy ϵ_i with degeneracy factor g_i . If n_i is the average number of particles in i -th level, then calculate the number of distinct microstates corresponding to the set of occupation numbers $\{n_i\}$.
- b) Eleven (11) particles are to be distributed in nine (9) degenerate energy levels with energy ϵ . Calculate the number of ways of distributing these particles if they are (i) classical identical particles, (ii) Fermions and (iii) Bosons.
- $4+(2+2+2)=10$
